


Biochemistry of vitamins-1.

Fat soluble vitamins

Lecture #24

Lecturer: Alexander Koval

Plan of lectures, part 2



Introduction to Vitamins




- Vitamins are organic molecules that function in a wide variety of capacities within the body.
 - The most prominent function is as cofactors for enzymatic reactions.
 - The distinguishing feature of the vitamins is that they generally cannot be synthesized by mammalian cells and, therefore, must be supplied in the diet.

The Vitamins are of Two Distinct Types:

Lipid-Soluble	Water-Soluble
<ol style="list-style-type: none">1. Vitamin A2. Vitamin D3. Vitamin E4. Vitamin K	<ol style="list-style-type: none">1. Thiamin (B₁)2. Riboflavin (B₂)3. Niacin (B₅)4. Pantothenic Acid (B₃)5. Pyridoxine (Pyridoxal, Pyridoxol, Pyridoxamine) (B₆)6. Biotin (H)7. Cobalamin (B₁₂)8. Folic Acid (B₉)9. Ascorbic Acid (C)

Lipid-Soluble Vitamins


* Adult daily requirement



Lipid-Soluble Vitamins (cont'd)

* Adult daily requirement

Provitamin	Functional form	Important for
------------	-----------------	---------------




Tocopherols

10
Cereals
Liver
Eggs
Seed oil




Tocopherols

Reducing agent




Anti-oxidant



Phylloquinones

0.08
Intestinal
bacteria
Vegetables
Liver






Phyllohydroquinones



Blood clotting
(carboxylation
of plasma
proteins)




Vitamin A




- Vitamin A consists of three biologically active molecules, **retinol**, **retinal (retinaldehyde)** and **retinoic acid**.

Provitamin Form of Vitamin A



- Each of these compounds are derived from the plant precursor molecule, β -**carotene** (a member of a family of molecules known as **carotenoids**).
- β -carotene, which consists of two molecules of retinal linked at their aldehyde ends, is also referred to as the provitamin form of vitamin A.

Vitamin A1 and its precursor and derivatives



Transport in the blood


- Ingested β -carotene is cleaved in the lumen of the intestine by **β -carotene dioxygenase** to yield retinal.
- Retinal is reduced to retinol by **retinaldehyde reductase**, (NADPH enzyme) within the intestines.
- Retinol is esterified to *palmitic acid* and delivered to the blood via **chylomicrons**.
 - The uptake of chylomicron remnants by the liver results in delivery of retinol to this organ for storage as a lipid ester within lipocytes.
- Transport of retinol from the liver to extrahepatic tissues occurs by binding of hydrolyzed retinol to aporetinol binding protein (RBP). the retinol-RBP complex is then transported to the cell surface within the Golgi and secreted.
 - Within extrahepatic tissues retinol is bound to cellular retinol binding protein (CRBP).
 - Plasma transport of retinoic acid is accomplished by binding to albumin.

Gene Control Exerted by Retinol and Retinoic Acid

- Within cells both retinol and retinoic acid bind to specific receptor proteins. Following binding, the receptor-vitamin complex interacts with specific sequences in several genes involved in growth and differentiation and affects expression of these genes.
- In this capacity retinol and retinoic acid are considered hormones of the steroid/thyroid hormone superfamily.
 - Vitamin D also acts in a similar capacity.
- Several genes whose patterns of expression are altered by retinoic acid are involved in the earliest processes of embryogenesis including the differentiation of the three germ layers, organogenesis and limb development.

Vision and the Role of Vitamin A

- Photoreception in the eye is the function of two specialized cell types located in the retina; the rod and cone cells.




- Both rod and cone cells contain a photoreceptor pigment in their membranes. The photosensitive compound of most mammalian eyes is a protein called **opsin** to which is covalently coupled an **aldehyde of vitamin A**.

Rhodopsin


- The opsin of rod cells is called **scotopsin**.
- The photoreceptor of rod cells is specifically called **rhodopsin** or **visual purple**. This compound is a complex between **scotopsin** and the **11-cis-retinal** (also called **11-cis-retinene**) form of vitamin A.
- Rhodopsin is a serpentine receptor imbedded in the membrane of the rod cell.
- Coupling of 11-cis-retinal occurs at three of the transmembrane domains of rhodopsin.
- Intracellularly, rhodopsin is coupled to a specific **G-protein** called **transducin**.

Processes in Retina

- When the rhodopsin is exposed to light it is bleached releasing the **11-cis-retinal** from opsin.
- Absorption of photons by 11-cis-retinal triggers a series of conformational changes on the way to conversion **all-trans-retinal**.
 - One important conformational intermediate is **metarhodopsin II**.



The double bond between C-11 and C-12 is isomerized from cis to trans by light



all-trans-Retinal

Vision Act

- The release of opsin results in a conformational change in the photoreceptor.
 - This conformational change activates transducin, leading to an increased GTP-binding by the α -subunit of transducin.
 - Binding of GTP releases the α -subunit from the inhibitory β - and γ -subunits.
 - The GTP-activated α -subunit in turn activates an associated phosphodiesterase; an enzyme that hydrolyzes cyclic-GMP (cGMP) to GMP. Cyclic GMP is required to maintain the Na^+ channels of the rod cell in the open conformation.
 - The drop in cGMP concentration results in complete closure of the Na^+ channels.
 - Metarhodopsin II appears to be responsible for initiating the closure of the channels. The closing of the channels leads to hyperpolarization of the rod cell with concomitant propagation of nerve impulses to the brain.

Additional Role of Retinol

- Retinol also functions in the synthesis of certain glycoproteins and mucopolysaccharides necessary for mucous production and normal growth regulation.
- This is accomplished by phosphorylation of retinol to **retinyl phosphate** which then functions similarly to **dolichol phosphate**.


Clinical Significances of Vitamin A Deficiency

- Vitamin A is stored in the liver and deficiency of the vitamin occurs only after prolonged lack of dietary intake.
 - The earliest symptoms of vitamin A deficiency are **night blindness**.
 - Additional early symptoms include **follicular hyperkeratinosis**, increased **susceptibility to infection** and **cancer** and **anemia** equivalent to iron deficient anemia.
 - Prolonged lack of vitamin A leads to deterioration of the eye tissue through progressive keratinization of the cornea, a condition known as **xerophthalmia**.

Antioxidant and toxicity


- The increased risk of cancer in vitamin deficiency is thought to be the result of a depletion in β -carotene.
- Beta-carotene is a very effective **antioxidant** and is suspected to reduce the risk of cancers known to be initiated by the production of free radicals.
 - Of particular interest is the potential benefit of increased β -carotene intake to reduce the risk of lung cancer in smokers.
 - However, caution needs to be taken when increasing the intake of any of the lipid soluble vitamins.
 - Excess accumulation of vitamin A in the liver can lead to **toxicity** which manifests as **bone pain, hepatosplenomegaly, nausea** and **diarrhea**.

Vitamin D



- Active calcitriol is derived from **ergosterol** (produced in plants) and from **7-dehydrocholesterol** (produced in the skin).
- **Ergocalciferol** (vitamin D₂) is formed by uv irradiation of ergosterol.

D2 & D3





- In the skin **7-dehydrocholesterol** is converted to **cholecalciferol** (vitamin D₃) following uv irradiation.
- Vitamin D₂ and D₃ are processed to D₂-calcitriol and D₃-calcitriol, respectively, by the same enzymatic pathways in the body.

Hydroxylation


- Cholecalciferol (or ergocalciferol) are absorbed from the intestine and transported to the liver bound to a specific **vitamin D-binding protein**.
- In the liver cholecalciferol is hydroxylated at the 25 position by a specific **D₃-25-hydroxylase** generating 25-hydroxy-D₃ [25-(OH)D₃] which is the major circulating form of vitamin D.
- Conversion of 25-(OH)D₃ to its biologically active form, calcitriol, occurs through the activity of a specific **D₃-1-hydroxylase** present in the proximal convoluted tubules of the kidneys, and in bone and placenta.
- 25-(OH)D₃ can also be hydroxylated at the 24 position by a specific **D₃-24-hydroxylase** in the kidneys, intestine, placenta and cartilage.

Metabolism of vitamin D





25-hydroxyvitamin D₃



1,25-dihydroxyvitamin D₃


Calcitriol Functions

- Calcitriol functions in concert with **parathyroid hormone (PTH)** and calcitonin to regulate serum calcium and phosphorous levels.
- PTH is released in response to low serum calcium and induces the production of calcitriol.
- In contrast, reduced levels of PTH stimulate synthesis of the inactive $24,25\text{-(OH)}_2\text{D}_3$.
- In the intestinal epithelium, calcitriol functions as a steroid hormone in inducing the expression of calbindin $\text{D}_{28\text{K}}$, a protein involved in intestinal calcium absorption.

Calcium and Phosphate Metabolism

- The increased absorption of calcium ions requires concomitant absorption of a negatively charged counter ion to maintain electrical neutrality.
 - The predominant counter ion is P_i .
 - When plasma calcium levels fall the major sites of action of calcitriol and PTH are bone where they stimulate bone resorption and the kidneys where they inhibit calcium excretion by stimulating reabsorption by the distal tubules.
 - The role of calcitonin in calcium homeostasis is to decrease elevated serum calcium levels by inhibiting bone resorption.

Clinical Significance of Vitamin D Deficiency



- As a result of the addition of vitamin D to milk, deficiencies in this vitamin are rare in this country.
- The main symptom of vitamin D deficiency in children is **rickets** and in adults is **osteomalacia**.

Rickets



Before vitamin D treatment





After 14 months of vitamin D treatment

- **Rickets** is characterized improper mineralization during the development of the bones resulting in soft bones.
- **Osteomalacia** is characterized by demineralization of previously formed bone leading to increased softness and susceptibility to fracture.


Functions of Vitamin E

- The major function of vitamin E is to act as a natural antioxidant by scavenging free radicals and molecular oxygen.
 - In particular vitamin E is important for preventing peroxidation of polyunsaturated membrane fatty acids.
 - The vitamins E and C are interrelated in their antioxidant capabilities. Active α -tocopherol can be regenerated by interaction with vitamin C following scavenge of a peroxy free radical.
 - Alternatively, α -tocopherol can scavenge two peroxy free radicals and then be conjugated to glucuronate for excretion in the bile.

Vitamin K




- The K vitamins exist naturally as K₁ (**phylloquinone**) in green vegetables and K₂ (**menaquinone**) produced by intestinal bacteria and K₃ is synthetic **menadione**.
- When administered, vitamin K₃ is alkylated to one of the vitamin K₂ forms of menaquinone.



Function of Vitamins K

- The major function of the K vitamins is in the maintenance of normal levels of the blood clotting proteins, **factors II, VII, IX, X** and **protein C** and **protein S**, which are synthesized in the liver as inactive precursor proteins.
- Conversion from inactive to active clotting factor requires a posttranslational modification of specific glutamate (E) residues. This modification is a carboxylation and the enzyme responsible requires vitamin K as a cofactor.
- The resultant modified E residues are **γ -carboxyglutamate (gla)**.
 - This process is most clearly understood for factor II, also called **preprothrombin**. Prothrombin is modified preprothrombin.
 - The gla residues are effective calcium ion chelators.
 - Upon chelation of calcium, prothrombin interacts with phospholipids in membranes and is proteolysed to thrombin through the action of activated **factor X (Xa)**.

Conversion of Vitamins K



- During the carboxylation reaction reduced hydroquinone form of vitamin K is converted to a 2,3-epoxide form.
- The regeneration of the hydroquinone form requires an uncharacterized reductase.
- This latter reaction is the site of action of the **dicumarol** based anticoagulants such as **warfarin**.
- 1 monooxygenase;
2 carboxylase;
3 2,3-epoxide reductase
4 reductase
(by Murray et. al., 1996)

Clinical significance of Vitamin K Deficiency

- Naturally occurring vitamin K is absorbed from the intestines only in the presence of bile salts and other lipids through interaction with chylomicrons.
 - Therefore, fat malabsorptive diseases can result in vitamin K deficiency.
- The synthetic vitamin K₃ is water soluble and absorbed irrespective of the presence of intestinal lipids and bile.
- Since the vitamin K₂ form is synthesized by intestinal bacteria, deficiency of the vitamin in adults is rare.
 - However, long term antibiotic treatment can lead to deficiency in adults.
- The intestine of newborn infants is sterile, therefore, vitamin K deficiency in infants is possible if lacking from the early diet.
- The primary symptom of a deficiency in infants is a **hemorrhagic syndrome**.


Biochemistry of vitamins-2.

Water soluble vitamins

Lecture #25


Lecturer: Alexander Koval

Thiamin



- Thiamin is also known as **vitamin B₁**.
 - Thiamin is derived from a substituted pyrimidine and a thiazole which are coupled by a methylene bridge.
- Thiamin is rapidly converted to its active form, **thiamin pyrophosphate, TPP**, in the brain and liver by a specific enzymes, *thiamin diphosphotransferase*.

Thiamin Pyrophosphate



- TPP is necessary as a cofactor for the **pyruvate** and **α -ketoglutarate dehydrogenase** catalyzed reactions as well as the **transketolase** catalyzed reactions of the pentose phosphate pathway.
 - A deficiency in thiamin intake leads to a severely reduced capacity of cells to generate energy as a result of its role in these reactions.

Dietary Requirement

- The dietary requirement for thiamin is proportional to the caloric intake of the diet and ranges from 1.0 - 1.5 mg/day for normal adults.
- If the carbohydrate content of the diet is excessive then an in thiamin intake will be required.

Clinical Significances of Thiamin Deficiency


- The earliest symptoms of thiamin deficiency include constipation, appetite suppression, nausea as well as mental depression, peripheral neuropathy and fatigue.
 - Chronic thiamin deficiency leads to more severe neurological symptoms including ataxia, mental confusion and loss of eye coordination.
 - Other clinical symptoms of prolonged thiamin deficiency are related to cardiovascular and musculature defects.

Beriberi

- The severe thiamin deficiency disease known as **Beriberi**, is the result of a diet that is carbohydrate rich and thiamin deficient.
- An additional thiamin deficiency related disease is known as **Wernicke-Korsakoff** syndrome.
 - This disease is most commonly found in chronic alcoholics due to their poor dietic lifestyles.

Riboflavin

- Riboflavin is also known as **vitamin B₂**.




Riboflavin

FMN & FAD

- Riboflavin is the precursor for the coenzymes, **flavin mononucleotide (FMN)** and **flavin adenine dinucleotide (FAD)**.
 - The enzymes that require FMN or FAD as cofactors are termed flavoproteins.
 - Several flavoproteins also contain metal ions and are termed metalloflavoproteins.
 - Both classes of enzymes are involved in a wide range of redox reactions, e.g. *succinate dehydrogenase* and *xanthine oxidase*.
- During the course of the enzymatic reactions involving the flavoproteins the reduced forms of FMN and FAD are formed, FMNH_2 and FADH_2 , respectively.

Structure of FAD




- nitrogens 1 & 5 carry hydrogens in FADH₂


Clinical Significances of Flavin Deficiency

- Riboflavin deficiencies are rare in humans due to the presence of adequate amounts of the vitamin in eggs, milk, meat and cereals.
 - Riboflavin deficiency is often seen in chronic alcoholics due to their poor dietetic habits.
- Symptoms associated with riboflavin deficiency include, glossitis, seborrhea, angular stomatitis, cheilosis and photophobia.
 - Riboflavin decomposes when exposed to visible light.
 - This characteristic can lead to riboflavin deficiencies in newborns treated for hyperbilirubinemia by phototherapy.

Niacin



Nicotinic Acid




Nicotinamide

Vitamin B₅, NAD⁺, NADP⁺

- Niacin (nicotinic acid and nicotinamide) is also known as vitamin B₅.
 - Both nicotinic acid and nicotinamide can serve as the dietary source of vitamin B₅.
 - Niacin is required for the synthesis of the active forms of vitamin B₅, nicotinamide adenine dinucleotide (NAD⁺) and nicotinamide adenine dinucleotide phosphate (NADP⁺).
 - Both NAD⁺ and NADP⁺ function as cofactors for numerous dehydrogenase, e.g., lactate and malate dehydrogenases.

Structure of NAD⁺



- NADH is shown in the box insert.
- The -OH phosphorylated in NADP⁺ is indicated by the arrow.

Daily Requirement

- Niacin is not a true vitamin in the strictest definition since it can be derived from the amino acid tryptophan.
 - However, the ability to utilize tryptophan for niacin synthesis is inefficient (60 mg of tryptophan are required to synthesize 1 mg of niacin).
 - Also, synthesis of niacin from tryptophan requires vitamins B₁, B₂ and B₆ which would be limiting in themselves on a marginal diet.
- The recommended daily requirement for niacin is 13 - 19 niacin equivalents (NE) per day for a normal adult.
- One NE is equivalent to 1 mg of free niacin).


Clinical Significances of Niacin and Nicotinic Acid

- A diet deficient in niacin (as well as tryptophan) leads to glossitis of the tongue, dermatitis, weight loss, diarrhea, depression and dementia.
 - The severe symptoms, depression, dermatitis and diarrhea, are associated with the condition known as **pellagra**.
- Several physiological conditions (e.g. Hartnup disease and **malignant carcinoid syndrome**) as well as certain drug therapies (e.g. isoniazid) can lead to niacin deficiency.
 - In Hartnup disease tryptophan absorption is impaired and in malignant carcinoid syndrome tryptophan metabolism is altered resulting in excess serotonin synthesis.
 - Isoniazid (the hydrazide derivative of isonicotinic acid) is the primary drug for chemotherapy of tuberculosis.

Therapeutic use


- Nicotinic acid (but not nicotinamide) when administered in pharmacological doses of 2 - 4 g/day lowers plasma cholesterol levels and has been shown to be a useful therapeutic for hypercholesterolemia.
 - The major action of nicotinic acid in this capacity is a reduction in fatty acid mobilization from adipose tissue.
 - Although nicotinic acid therapy lowers blood cholesterol it also causes a depletion of glycogen stores and fat reserves in skeletal and cardiac muscle.
 - Additionally, there is an elevation in blood glucose and uric acid production.
 - For these reasons nicotinic acid therapy is not recommended for diabetics or persons who suffer from gout.

Pantothenic Acid



- Pantothenic acid is also known as vitamin B₃.
 - Pantothenic acid is formed from **β-alanine** and **pantoic acid**.

Coenzyme A




- Pantothenate is required for synthesis of **coenzyme A, CoA** and is a component of the **acyl carrier protein (ACP)** domain of **fatty acid synthase**.
 - Pantothenate is, therefore, required for the metabolism of carbohydrate via the TCA cycle and all fats and proteins.


Requirements and Deficiency

- At least **70 enzymes** have been identified as requiring CoA or ACP derivatives for their function.
- Deficiency of pantothenic acid is extremely rare due to its widespread distribution in **whole grain cereals, legumes and meat**.
 - Symptoms of pantothenate deficiency are difficult to assess since they are subtle and resemble those of **other B vitamin deficiencies**.


Vitamin B₆



pyridoxol




pyridoxal



pyridoxamine

- **Pyridoxal, pyridoxamine and pyridoxol (pyridoxine)** are collectively known as vitamin B₆ (**pyridoxine**).

Pyridoxal Phosphate



pyridoxalphosphate
(coenzyme)

- All three compounds are efficiently converted to the biologically active form of vitamin B₆, **pyridoxal phosphate**.
 - This conversion is catalyzed by the ATP requiring enzyme, pyridoxal kinase.


Requirement of B₆

- **Pyridoxal phosphate** functions as a cofactor in enzymes involved in transamination reactions required for the synthesis and catabolism of the amino acids as well as in glycogenolysis as a cofactor for glycogen phosphorylase.
- The requirement for vitamin B₆ in the diet is proportional to the level of protein consumption ranging from 1.4 - 2.0 mg/day for a normal adult.
 - During pregnancy and lactation the requirement for vitamin B6 increases approximately 0.6 mg/day.

Deficiencies of vitamin B₆

- Deficiencies of vitamin B₆ are rare and usually are related to an overall deficiency of all the B-complex vitamins.
- **Isoniazid** and **penicillamine** (used to treat rheumatoid arthritis and cystinurias) are two drugs that complex with pyridoxal and pyridoxal phosphate resulting in a deficiency in this vitamin.

Biotin




- **Biotin** is the cofactor required of enzymes that are involved in carboxylation reactions, e.g. **acetyl-CoA carboxylase** and **pyruvate carboxylase**.

Biotin Functions

- Biotin is found in numerous foods and also is synthesized by intestinal bacteria and as such deficiencies of the vitamin are rare.
- Deficiencies are generally seen only after long antibiotic therapies which deplete the intestinal fauna or following excessive consumption of raw eggs.
- The latter is due to the affinity of the egg white protein, avidin, for biotin preventing intestinal absorption of the biotin.

Vitamin B₁₂, Cobalamin



- **Cobalamin** is more commonly known as **vitamin B₁₂**.
- Vitamin B₁₂ is composed of a complex **tetrapyrrole ring structure (corrin ring)** and a **cobalt** ion in the center.

Synthesis of B₁₂

- Vitamin B₁₂ is synthesized exclusively by microorganisms and is found in the liver of animals bound to protein as **methycobalamin** or **5'-deoxyadenosylcobalamin**.
- The vitamin must be hydrolyzed from protein in order to be active.

Vitamin B₁₂ Digestion, Transportation

- Hydrolysis occurs in the stomach by gastric acids or the intestines by trypsin digestion following consumption of animal meat. The vitamin is then bound by intrinsic factor, a protein secreted by parietal cells of the stomach, and carried to the ileum where it is absorbed.
- Following absorption the vitamin is transported to the liver in the blood bound to transcobalamin II.

Significance of vitamin B₁₂

- There are only two clinically significant reactions in the body that require vitamin B₁₂ as a cofactor.
- During the catabolism of fatty acids with an odd number of carbon atoms and the amino acids *valine*, *isoleucine* and *threonine* the resultant **propionyl-CoA** is converted to **succinyl-CoA** for oxidation in the TCA cycle.
- One of the enzymes in this pathway, methylmalonyl-CoA mutase, requires vitamin B₁₂ as a cofactor in the conversion of methylmalonyl-CoA to succinyl-CoA.
- The 5'-deoxyadenosine derivative of cobalamin is required for this reaction.


The Role in Reactions of Conversion

- The second reaction requiring vitamin B12 catalyzes the conversion of *homocysteine* to *methionine* and is catalyzed by **methionine synthase**.
- This reaction results in the transfer of the methyl group from **N⁵-methyltetrahydrofolate** to hydroxycobalamin generating **tetrahydrofolate (THF)** and **methylcobalamin** during the process of the conversion.

Clinical Significances of B₁₂ Deficiency

- The liver can store up to six years worth of vitamin B₁₂, hence deficiencies in this vitamin are rare.
- **Pernicious anemia** is a **megaloblastic anemia** resulting from vitamin B₁₂ deficiency that develops as a result a lack of intrinsic factor in the stomach leading to malabsorption of the vitamin.
 - The anemia results from impaired DNA synthesis due to a block in purine and thymidine biosynthesis. The block in nucleotide biosynthesis is a consequence of the effect of vitamin B₁₂ on folate metabolism.
 - When vitamin B₁₂ is deficient essentially all of the folate becomes trapped as the N⁵-methylTHF derivative as a result of the loss of functional methionine synthase.
- This trapping prevents the synthesis of other THF derivatives required for the purine and thymidine nucleotide biosynthesis pathways.

Folic Acid








positions 7 & 8 carry hydrogens in dihydrofolate (DHF)
positions 5-8 carry hydrogens in tetrahydrofolate (THF)

Folic acid is a conjugated molecule consisting of a **pteridine** ring structure linked to **para-aminobenzoic acid (PABA)** that forms **pteroic acid**. Folic acid itself is then generated through the conjugation of **glutamic acid** residues to **pteroic acid**.

- Folic acid is obtained primarily from *yeasts* and *leafy vegetables* as well as animal *liver*.
 - Animal cannot synthesize PABA nor attach glutamate residues to pteronic acid, thus, requiring folate intake in the diet.
- When stored in the liver or ingested folic acid exists in a **polyglutamate form**.
 - Intestinal mucosal cells remove some of the glutamate residues through the action of the lysosomal enzyme, **conjugase**. The removal of glutamate residues makes folate less negatively charged (from the polyglutamic acids) and therefore more capable of passing through the basal laminal membrane of the epithelial cells of the intestine and into the bloodstream.
- Folic acid is reduced within cells (principally the liver where it is stored) to tetrahydrofolate (THF also H₄folate) through the action of dihydrofolate reductase (DHFR), an NADPH-requiring enzyme.

Function of THF



- The function of THF derivatives is to carry and transfer various forms of one carbon units during biosynthetic reactions. The one carbon units are either methyl, methylene, methenyl, formyl or formimino groups.
- Active center of tetrahydrofolate (THF). Note that the N⁵ position is the site of attachment of methyl groups, the N¹⁰ the site for attachment of formyl and formimino groups and that both N⁵ and N¹⁰ bridge the methylene and methenyl groups.

Role of Folic Acid in Metabolism

- These one carbon transfer reactions are required in the biosynthesis of serine, methionine, glycine, choline and the purine nucleotides and dTMP.
- The ability to acquire choline and amino acids from the diet and to salvage the purine nucleotides makes the role of N^5,N^{10} -methylene-THF in dTMP synthesis the most metabolically significant function for this vitamin.
 - The role of vitamin B_{12} and N^5 -methyl-THF in the conversion of homocysteine to methionine also can have a significant impact on the ability of cells to regenerate needed THF.


Clinical Significance of Folate Deficiency

- Folate deficiency results in complications nearly identical to those described for vitamin B₁₂ deficiency.
- The most pronounced effect of folate deficiency on cellular processes is upon DNA synthesis.
 - This is due to an impairment in dTMP synthesis which leads to cell cycle arrest in S-phase of rapidly proliferating cells, in particular hematopoietic cells.
 - The result is **megaloblastic anemia** as for vitamin B₁₂ deficiency.
 - The inability to synthesize DNA during erythrocyte maturation leads to abnormally large erythrocytes termed **macrocytic anemia**.

Cases of Folate Deficiencies

- Folate deficiencies are rare due to the adequate presence of folate in food.
- Poor dietary habits as those of chronic alcoholics can lead to folate deficiency.
- The predominant causes of folate deficiency in non-alcoholics are impaired absorption or metabolism or an increased demand for the vitamin.
- The predominant condition requiring an increase in the daily intake of folate is pregnancy. This is due to an increased number of rapidly proliferating cells present in the blood. The need for folate will nearly double by the third trimester of pregnancy.
- Certain drugs such as anticonvulsants and oral contraceptives can impair the absorption of folate.
 - Anticonvulsants also increase the rate of folate metabolism.

Ascorbic Acid




- Ascorbic acid is more commonly known as **vitamin C**.
 - Ascorbic acid is derived from glucose via the **uronic acid pathway**.
 - The enzyme **L-gulonolactone oxidase** responsible for the conversion of gulonolactone to ascorbic acid is absent in primates making ascorbic acid required in the diet.

Functions of Ascorbate

- The active form of vitamin C is ascorbic acid itself.
- The main function of ascorbate is as a reducing agent in a number of different reactions.
 - Vitamin C has the potential to reduce cytochromes a and c of the respiratory chain as well as molecular oxygen.
 - Vitamin C also is necessary for bone remodeling due to the presence of collagen in the organic matrix of bones.

Proline Hydroxylation



- The most important reaction requiring ascorbate as a cofactor is the hydroxylation of proline residues in collagen.
- Vitamin C is, therefore, required for the maintenance of normal connective tissue as well as for wound healing since synthesis of connective tissue is the first event in wound tissue remodeling.

Vitamin C as a Cofactor

- Several other metabolic reactions require vitamin C as a cofactor.
 - These include the catabolism of **tyrosine** and the synthesis of **epinephrine** from tyrosine and the synthesis of the **bile acids**.
 - It is also believed that vitamin C is involved in the process of **steroidogenesis** since the adrenal cortex contains high levels of vitamin C which are depleted upon adrenocorticotrophic hormone (ACTH) stimulation of the gland.

Vitamin C Deficiency

- Deficiency in vitamin C leads to the disease **scurvy** due to the role of the vitamin in the post-translational modification of collagens.
 - Scurvy is characterized by easily bruised skin, muscle fatigue, soft swollen gums, decreased wound healing and hemorrhaging, osteoporosis, and anemia.
 - Vitamin C is readily absorbed and so the primary cause of vitamin C deficiency is poor diet and/or an increased requirement.
 - The primary physiological state leading to an increased requirement for vitamin C is severe stress (or trauma). This is due to a rapid depletion in the adrenal stores of the vitamin.
- The reason for the decrease in adrenal vitamin C levels is unclear but may be due either to redistribution of the vitamin to areas that need it or an overall increased utilization.

The End

Thank you
for your attention